

Exercises on General Relativity TVI TMP-TC1

Problem set 7 extra, due December 8th

This problem sheet is discussed instead of the lecture.

Exercise 1 – Small exercises

- (i) Compute the mass dimension of the scalar field ϕ and constants λ_n of the following action in eight space-time dimensions:

$$S = \int d^8x \left(-\frac{1}{2} \phi \square \phi - \sum_{n=2}^4 \frac{\lambda_n}{n!} \phi^n \right).$$

- (ii) Write the following tensors in a local coordinate basis using the basis theorem: (a) metric, (b) inverse metric, and (c) rank-(2,2) tensor T .
- (iii) Calculate the 1-forms dh and dk , where

$$h(x, y, z) = x^2 y^2 + x^3 z, \quad k(x, y) = \frac{1}{\sqrt{x^2 + y^2}}.$$

- (iv) Solve for the unknown vector X^α in the following equation:

$$\epsilon_{\alpha\beta\gamma} X^\beta A^\gamma = B_\alpha \quad \text{and} \quad X^\alpha C_\alpha = k, \quad \text{where} \quad A^\alpha B_\alpha = 0 \quad (\text{in } \mathbb{R}^3) \quad (1)$$

where k is a non-zero scalar, A, B and C are linearly independent vectors and ϵ is the 3D totally antisymmetric tensor.

Exercise 2 – Working on a sphere

- (i) Reconsider the stereographic projections of an n -sphere of problem set 7 for $n = 2$. The coordinates on \mathbb{S}^2 we want to use here is given by the stereographic projection from the north pole $N := (0, 0, 1)$:

$$y_1 = \frac{x_1}{1-z}, \quad y_2 = \frac{x_2}{1-z} \quad (2)$$

Let the vector fields X and Y on $U^+ := \mathbb{S}^2 \setminus \{N\}$ be defined in these coordinates by

$$X = y_2 \frac{\partial}{\partial y_1} - y_1 \frac{\partial}{\partial y_2}, \quad Y = y_1 \frac{\partial}{\partial y_1} + y_2 \frac{\partial}{\partial y_2}. \quad (3)$$

Express these two vector fields in coordinates corresponding to the stereographic projection from the south pole $S := (0, 0, -1)$.

- (ii) Consider a surface $S \subseteq \mathbb{R}^3$ parametrized by a differentiable map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^3, f(u^1, u^2)$. Let $g(x, y) = \sqrt{R^2 - x^2 - y^2}$ with a fixed real positive number R . The graph of f can be parametrised as follows:

$$f = (u^1, u^2, g(u^1, u^2)). \quad (4)$$

This provides a parametrisation of the (open) northern hemisphere.

Judge whether this is a regular parametrisation of the surface S , i.e. $\frac{\partial f}{\partial u^1}$ and $\frac{\partial f}{\partial u^2}$ are linear independent.